
COOK COUNTY DEPARTMENT OF PUBLIC HEALTH
Influenza A (H1N1) Vaccine Post-test Training

Direction: Circle the correct answer.

1. There are four (4) H1N1 types of vaccine manufactured for use in the U.S.
 - a. True
 - b. False

2. The Live Attenuated Intranasal Vaccine (LAIV-H1N1) is a weakened strain of the H1N1 virus and is intended for healthy people ages 2 – 49 years of age.
 - a. True
 - b. False

3. Active inhalation (i.e., sniffing) is required during administration of LAIV-H1N1.
 - a. True
 - b. False

4. LAIV-H1N1 must be kept frozen until used.
 - a. True
 - b. False

5. There is no preservative in the single dose preparations of H1N1 vaccine licensed for use in children < 4 years of age.
 - a. True
 - b. False

6. If a student has received an Influenza A (H1N1) vaccination in the past 28 days he/she will not be vaccinated by CCDPH.
 - a. True
 - b. False

7. Immunocompromised people should not receive LAIV-H1N1.
 - a. True
 - b. False

8. If the person to be vaccinated has an illness with fever > 100° F or other active infection they will be vaccinated by CCDPH.
 - a. True
 - b. False

9. Hypersensitivity to eggs or any component of the vaccine and/or a life threatening reaction to previous influenza vaccine is contraindications to receiving vaccination.
 - a. True
 - b. False

10. Pregnant people, people younger than 2 years of age, people over the age of 50 and people with a history of asthma or active wheezing will not receive LAIV-H1N1.
 - a. True
 - b. False

11. If the person to be vaccinated has received live virus vaccines like FluMist, MMR, or Varicella in the past 28 days they should not be vaccinated with LAIV-H1N1.
 - a. True
 - b. False

12. People with a weakened immune system which would make him/her more likely to contract an infection should be vaccinated with LAIV-H1N1.
 - a. True
 - b. False

13. People < 18 years of age receiving Aspirin-containing therapy should receive LAIV-H1N1.
 - a. True
 - b. False

14. People allergic to gentamicin, gelatin, or arginine should not receive LAIV-H1N1.
 - a. True
 - b. False

15. Antiviral medications taken 48 hours prior to vaccination can lower or prevent the vaccinated person from responding to the vaccine and the person may not get immune protection from the vaccine.
 - a. True
 - b. False

16. People 6 months of age and older and people < 10 years of age need two (2) doses of H1N1 vaccine to confer immunity.
 - a. True
 - b. False

17. People with a history of Guillain-Barre´ syndrome should receive H1N1 vaccine.
 - a. True
 - b. False