

**This is an official**  
**CDC Health Advisory**

\* Please acknowledge receipt of this HAN by following the link at the bottom of the advisory.

Distributed via Health Alert Network  
August 6, 2009, 19:00 ET (7:00 PM ET)  
CDCHAN-00297-09-08-06-ADV-N

**CDC Updates Recommendations for the Amount of Time Persons with  
Influenza-Like Illness should be Away from Others**

On August 5, 2009, CDC changed its recommendation related to the amount of time people with influenza-like illness should stay away from others (the exclusion period). New guidance indicates that people with influenza-like illness should stay home for at least 24 hours after their fever is gone (without the use of fever-reducing medicine). A fever is defined as having a temperature of 100° Fahrenheit or 37.8° Celsius or greater.

This is a change from the previous recommendation that ill persons stay home for 7 days after illness onset or until 24 hours after the resolution of symptoms, whichever was longer.

The new recommendation applies to camps, schools, businesses, mass gatherings, and other community settings where the majority of people are not at increased risk for influenza complications. CDC recommends this exclusion period regardless of whether or not antiviral medications are used. This guidance does not apply to health care settings where the exclusion period continues to be for 7 days from symptom onset or until 24 hours after the resolution of symptoms, whichever is longer. (See [http://www.cdc.gov/h1n1flu/guidelines\\_infection\\_control.htm](http://www.cdc.gov/h1n1flu/guidelines_infection_control.htm) for guidance on infection control in health care settings.)

Decisions about extending the exclusion period should be made at the community level, in conjunction with local and state health officials. More stringent guidelines and longer periods of exclusion – for example, until complete resolution of all symptoms – may be considered for people returning to a setting where high numbers of high-risk people may be exposed.

This exclusion period guidance for the community setting is based on epidemiologic data about the overall risk of severe illness and death. The new recommendation attempts to balance the risks of acquiring illness from influenza and the potential benefits of decreasing transmission through the exclusion of ill persons with the goal of minimizing social disruption. This guidance will continue to be updated as more information becomes available.

To read the complete revised guidance see: <http://www.cdc.gov/h1n1flu/guidance/exclusion.htm>

This change in our recommendation has affected content on a number of other pages, including the following:

Visit [http://www.cdc.gov/h1n1flu/guidance\\_homecare.htm](http://www.cdc.gov/h1n1flu/guidance_homecare.htm) for more information on caring for sick persons in the home.

Visit <http://www.cdc.gov/h1n1flu/qa.htm> questions and answers about H1N1 influenza.

For more general information on H1N1, go to <http://www.cdc.gov/h1n1flu/>.

A copy of the formatted HAN is attached in PDF form.

**Health Alert** conveys the highest level of importance; warrants immediate action or attention.  
**Health Advisory** provides important information for a specific incident or situation; may not require immediate action.  
**Health Update** provides updated information regarding an incident or situation; unlikely to require immediate action.

##This Message was distributed to State and Local Health Officers, Public Information Officers, Epidemiologists  
and HAN Coordinators as well as Clinician organizations##

=====  
You have received this message based upon the information contained within our emergency notification data base. If you have a different or  
additional e-mail or fax address that you would like us to use please contact your State-based Health Alert Network program at your State or local  
health department.  
=====